

PLANCHEREL-ROTACH ASYMPTOTICS FOR q -LAGUERRE ORTHOGONAL POLYNOMIALS WITH COMPLEX SCALING

RUIMING ZHANG

ABSTRACT. In this work we study the Plancherel-Rotach type asymptotics for q -Laguerre orthogonal polynomials with complex scaling. The main term of the asymptotics contains Ramanujan function $A_q(z)$ for the scaling parameter on the vertical line $\Re(s) = 2$, while the main term of the asymptotics involves the theta function for the scaling parameter in the strip $0 < \Re(s) < 2$. In the latter case the number theoretical property of the scaling parameter completely determines the order of the error term. These asymptotic formulas may provide insights to some new random matrix model and add a new link between special functions and number theory.

1. INTRODUCTION

The Plancherel-Rotach type asymptotics for classical orthogonal polynomials are essential to obtain universality results in random matrix theory. The associated random matrix models for q -orthogonal polynomials are still unknown today. It might be interesting to calculate the Plancherel-Rotach type asymptotics for q -orthogonal polynomials to gain some insights to the related random matrix models.

In [18] we studied Plancherel-Rotach type asymptotics for three families of q -orthogonal polynomials with real logarithmic scalings. They are Ismail-Masson polynomials $\{h_n(x|q)\}_{n=0}^{\infty}$, Stieltjes-Wigert polynomials $\{S_n(x; q)\}_{n=0}^{\infty}$ and q -Laguerre polynomials $\{L_n^{(\alpha)}(x; q)\}_{n=0}^{\infty}$. They are q -orthogonal polynomials associated with indeterminate moment problems. The asymptotics reveal a remarkable pattern which is quite different to the pattern associated with the classical Plancherel-Rotach asymptotics [24, 13]. The main term of asymptotics may contain Ramanujan function $A_q(z)$ or theta function according to the value of the scaling parameter.

In this paper, we derive the Plancherel-Rotach type asymptotics for q -Laguerre orthogonal polynomials under complex scaling. We also have explicit error bounds for the asymptotic formulas. When the scaling parameter s is in the vertical strip $0 < \Re(s) < 2$, the order of the error term is completely determined by the number theoretical property of the scaling parameter s .

In §2, we list some facts from q -series and number theory. We present the asymptotic formulas for q -Laguerre polynomials under complex scaling in §3. Finally we use a discrete analogue of Laplace method to derive these formulas in §4.

Through out this paper, We shall always assume that $0 < q < 1$ unless otherwise stated. We also assume that $s = \sigma + it$ is a complex number with $\sigma = \Re(s)$ and

1991 *Mathematics Subject Classification.* Primary 30E15. Secondary 33D45.

Key words and phrases. q -Orthogonal polynomials, Ramanujan function, q -Laguerre orthogonal polynomials, Plancherel-Rotach asymptotics, theta functions, complex scaling, random matrix.

$t = \Im(s)$. All the log and power functions are taken as their respective principle branches.

2. PRELIMINARIES

2.1. **q -series.** In this paper we follow the usual notation for q -series, [6, 9, 13]

$$(1) \quad (a; q)_0 := 1 \quad (a; q)_n := \prod_{k=0}^n (1 - aq^k), \quad \left[\begin{matrix} n \\ k \end{matrix} \right]_q := \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}},$$

and $n = \infty$ is allowed in the above definitions. Assume that $|z| < 1$, the q -Binomial theorem is [6, 9, 13]

$$(2) \quad \frac{(az; q)_\infty}{(z; q)_\infty} = \sum_{k=0}^{\infty} \frac{(a; q)_k}{(q; q)_k} z^k,$$

which defines an analytic function in the region $|z| < 1$. Its limiting case,

$$(3) \quad (z; q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q; q)_k} (-z)^k \quad z \in \mathbb{C}$$

is one of many q -exponential identities. Ramanujan function $A_q(z)$ is defined as [22, 13]

$$(4) \quad A_q(z) := \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q; q)_k} (-z)^k.$$

In order to express our error terms concisely, we also define a related function $B_q(z)$,

$$(5) \quad B_q(z) := \sum_{k=0}^{\infty} \frac{q^{k^2} z^k}{(q; q)_k}.$$

Clearly,

$$(6) \quad |A_q(z)| \leq B_q(|z|).$$

Since

$$k+1 \geq \frac{1 - q^{k+1}}{1 - q} \geq (k+1)q^k$$

and

$$\begin{aligned} B'_q(|z|) &= q \sum_{k=0}^{\infty} \frac{q^{k^2} |qz|^k (k+1) q^k}{(q; q)_k (1 - q^{k+1})} \\ &\leq \frac{q}{1 - q} B_q(|z|), \end{aligned}$$

thus,

$$(7) \quad |A'_q(z)| \leq B'_q(|z|) \leq \frac{q}{1 - q} B_q(|z|).$$

The theta function, [6, 9, 13]

$$(8) \quad \Theta(z|q) := \sum_{n=-\infty}^{\infty} q^{n^2} z^n$$

is clearly defined for any complex number $z \neq 0$. Jacobi triple product identity is

$$(9) \quad \Theta(z|q) = (q^2, -qz, -q/z; q^2)_\infty.$$

The q -Laguerre orthogonal polynomials $\left\{ L_n^{(\alpha)}(x; q) \right\}_{n=0}^\infty$ are defined as [13]

$$(10) \quad L_n^{(\alpha)}(x; q) = \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=0}^n \frac{q^{k^2+\alpha k} (-x)^k}{(q^{\alpha+1}; q)_k} \begin{bmatrix} n \\ k \end{bmatrix}_q,$$

for $\alpha > -1$. Let

$$(11) \quad s = \tau + 2 + i \frac{2\theta\pi}{\log q} \quad \tau, \theta \in \mathbb{R},$$

and

$$(12) \quad x_n(z, s) = zq^{-ns}.$$

Reverse the summation order to get

$$(13) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)}{(-zq^\alpha)^n q^{n^2(1-s)}} = \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{q^{k^2} e^{2nk\theta\pi i}}{(q^{\alpha+1}; q)_{n-k}} \left(-\frac{q^{\tau n}}{zq^\alpha} \right)^k.$$

For any real number x , then,

$$(14) \quad x = \lfloor x \rfloor + \{x\},$$

where the fractional part of x is $\{x\} \in [0, 1)$ and $\lfloor x \rfloor \in \mathbb{Z}$ is the greatest integer less or equal x . The arithmetic function

$$(15) \quad \chi(n) = 2 \left\{ \frac{n}{2} \right\} = n - 2 \left\lfloor \frac{n}{2} \right\rfloor,$$

which is the principal character modulo 2,

$$(16) \quad \chi(n) = \begin{cases} 1 & 2 \nmid n \\ 0 & 2 \mid n \end{cases}.$$

We will also make uses of the trivial inequality

$$(17) \quad |e^z - 1| \leq |z|e^{|z|}$$

for any $z \in \mathbb{C}$. The following lemma is from [18].

Lemma 2.1. *Given any $n \in \mathbb{N}$, if $a > 0$,*

$$(18) \quad \frac{(a; q)_\infty}{(a; q)_n} = (aq^n; q)_\infty = 1 + R_1(a; n)$$

with

$$(19) \quad |R_1(a; n)| \leq \frac{(-aq^2; q)_\infty aq^n}{1 - q},$$

while for $0 < aq < 1$,

$$(20) \quad \frac{(a; q)_n}{(a; q)_\infty} = \frac{1}{(aq^n; q)_\infty} = 1 + R_2(a; n)$$

with

$$(21) \quad |R_2(a; n)| \leq \frac{aq^n}{(1 - q)(aq; q)_\infty}.$$

Proof. It is clear from (3) that

$$R_1(a; n) = \sum_{k=1}^{\infty} \frac{q^{k(k-1)/2} (-aq^n)^k}{(q; q)_k} = -aq^n \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2} (-aq^n)^k}{(q; q)_{k+1}}.$$

Hence for $a > 0$, we have

$$\begin{aligned} \left| \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2} (-aq^n)^k}{(q; q)_{k+1}} \right| &\leq \frac{1}{1-q} \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q; q)_k} (aq^{n+1})^k \\ &\leq \frac{(-aq^2; q)_{\infty}}{1-q}, \end{aligned}$$

and (19) follows. Moreover from (2)

$$R_2(a; n) = aq^n \sum_{k=0}^{\infty} \frac{(aq^n)^k}{(q; q)_{k+1}},$$

hence

$$|R_2(a; n)| \leq \frac{aq^n}{(1-q)(aq^n; q)_{\infty}} \leq \frac{aq^n}{(1-q)(aq; q)_{\infty}},$$

which is (21) and the proof of the lemma is complete. $\square$

2.2. Generalized Irrational Measure. For an irrational number θ , Chebyshev's theorem implies that for any real number β , there exist infinitely many pairs of integers n and m with $n > 0$ such that [11]

$$(22) \quad n\theta = m + \beta + a_n \quad \text{with} \quad |a_n| \leq \frac{3}{n}.$$

Clearly, Chebyshev's theorem says that the arithmetic progression $\{n\theta\}_{n \in \mathbb{Z}}$ is ergodic in $\mathbb{R}$.

Definition 2.2. Given real numbers θ_j and β_j for $j = 1, \dots, N$, the generalized irrationality measure $\omega(\theta_1, \dots, \theta_N | \beta_1, \dots, \beta_N)$ of $\theta_1, \dots, \theta_N$ associated with $\beta_1, \dots, \beta_N$ is defined as the least upper bound of the set of real numbers r such that there exist infinitely many integers n and $m_1, \dots, m_N$ with $n > 0$ such that

$$(23) \quad |n\theta_j - \beta_j - m_j| < \frac{1}{n^{r-1}}$$

for $j = 1, \dots, N$.

Theorem 2.3. Let θ be an irrational number, then for any real number β , its generalized irrational measure associated with β is $\omega(\theta | \beta) \geq 2$.

Proof. This is a direct consequence of the Chebyshev's theorem. $\square$

The irrationality measure (or Liouville-Roth constant) $\mu(\theta)$ of a real number θ is defined as the least upper bound of the set of real numbers r such that [27]

$$(24) \quad 0 < |n\theta - m| < \frac{1}{n^{r-1}}$$

is satisfied by an infinite number of integer pairs (n, m) with $n > 0$. It is clear that we have

$$(25) \quad \omega(\theta|0) = \mu(\theta).$$

A real algebraic number θ of degree l if it is a root of an irreducible polynomial of degree l with integer coefficients. Liouville's theorem in number theory says that for a real algebraic number θ of degree l , there exists a positive constant $K(\theta)$ such that for any integer m and $n > 0$ we have

$$(26) \quad |n\theta - m| > \frac{K(\theta)}{n^{l-1}}.$$

A Liouville number is a real number θ such that for any positive integer l there exist infinitely many integers n and m with $n > 1$ such that [27]

$$(27) \quad 0 < |n\theta - m| < \frac{1}{n^{l-1}},$$

and the terms in the continued fraction expansion of every Liouville number are unbounded. Even though the set of all Liouville numbers is of Lebesgue measure zero, it is known that almost all real numbers are Liouville numbers topologically.

Theorem 2.4. *The generalized irrational measure $\omega(\theta|\beta)$ has the following properties:*

- (1) *For any real algebraic number θ of degree l , one has $\omega(\theta|0) \leq l$;*
- (2) *For a Liouville number θ , one has $\omega(\theta|0) = \infty$.*

Proof. The first assertion follows from the definition 2.2 and (26) while the last assertion follows directly from the definitions of the generalized irrational measure and the Liouville numbers. $\square$

It is clear that the quadratic irrationals θ such as $\sqrt{2}$ have $\omega(\theta|0) = 2$.

3. MAIN RESULTS

Given a real number θ , we define

$$(28) \quad \mathbb{S}(\theta) = \{\{n\theta\} : n \in \mathbb{N}\}.$$

When θ is a rational number, $\mathbb{S}(\theta)$ is a finite subset of $[0, 1)$. When θ is irrational, the set $\mathbb{S}(\theta)$ is a dense subset of $(0, 1)$, which is a consequence of Chebyshev's theorem.

Theorem 3.1. *Assume that $\alpha > -1$, given any nonzero complex number z , let s and $x_n(z, s)$ be defined as in (15) and (12), we have the following results for q -Laguerre polynomials:*

- (1) *When $\tau > 0$, we have*

$$(29) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} = 1 + r_{ql}(n|1)$$

with

$$(30) \quad |r_{ql}(n|1)| \leq \frac{q^{1-\alpha} B_q(q^{2-\alpha}|z|^{-1})}{(1-q)|z|} q^{\tau n}.$$

- (2) When $\tau = 0$ and θ is a rational number. For any $\lambda \in \mathbb{S}(\theta)$, there are infinitely many pair of integers n and m with $n > 0$ such that

$$(31) \quad n\theta = m + \lambda, \quad \lambda = \{n\theta\}.$$

For each of such integers,

$$(32) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} = A_q \left(\frac{e^{2\lambda\pi i}}{zq^\alpha} \right) + r_{ql}(n|2)$$

and the error is majorized as

$$(33) \quad |r_{ql}(n|2)| \leq \frac{7(-q^2; q)_\infty^2 B_q(|z|^{-1}q^{-\alpha})}{(1-q)^3(q; q)_\infty} \left\{ q^{n/2} + \frac{q^{n^2/4}}{(|z|q^\alpha)^{\lfloor n/2 \rfloor}} \right\}.$$

for n is sufficiently large.

- (3) When $\tau = 0$ and θ is an irrational number. Given any real number $\beta \in [0, 1)$ and a real number ρ with

$$(34) \quad 0 < \rho < \omega(\theta|\beta) - 1,$$

there are infinitely many pair of integers n and m with $n > 0$ such that

$$(35) \quad n\theta = m + \beta + \gamma_n \quad |\gamma_n| \leq \frac{1}{n^\rho}.$$

Then

$$(36) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} = A_q \left(\frac{e^{2\pi i\beta}}{zq^\alpha} \right) + e_{ql}(n|3)$$

with

$$(37) \quad |e_{ql}(n|3)| \leq \frac{48(-q^2; q)_\infty^2 B_q(|z|^{-1}q^{-\alpha})}{(1-q)^3(q; q)_\infty} \left\{ \frac{\log^2 n}{n^\rho} + \frac{q^{\nu_n^2}}{(|z|q^\alpha)^{\nu_n}} \right\}$$

and

$$(38) \quad \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor$$

for n is sufficiently large.

- (4) When $-2 < \tau < 0$ and both τ and θ are rational. Assume that for some $\lambda \in \mathbb{S}(-\tau)$ and some $\lambda_1 \in \mathbb{S}(\theta)$, there are infinite number of positive integers n such that

$$(39) \quad -n\tau = m + \lambda, \quad m = \lfloor -\tau n \rfloor, \quad \lambda = \{-\tau n\}$$

and

$$(40) \quad n\theta = m_1 + \lambda_1, \quad \lambda_1 = \{n\theta\}.$$

Then for each such n , m_1 , and m_2 , we have

$$(41) \quad \begin{aligned} L_n^{(\alpha)}(x_n(z, s); q) &= \frac{(-zq^\alpha)^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}}{(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}} \\ &\times \left\{ \Theta \left(-zq^\alpha q^{X(m) + \lambda} e^{-2\pi i\lambda_1} \mid q \right) + r_{ql}(n|4) \right\} \end{aligned}$$

with

$$(42) \quad |r_{ql}(n|4)| \leq \frac{30(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q})}{(1-q)^4(q; q)_\infty} \times \left\{ q^{\nu_n/2} + (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} \right\}$$

and

$$(43) \quad \nu_n = \min \left\{ \left\lfloor \frac{(2+\tau)n}{8} \right\rfloor, \left\lfloor \frac{-\tau n}{8} \right\rfloor \right\}.$$

for n sufficiently large.

- (5) When $-2 < \tau < 0$, τ is rational and θ is irrational. Given a real number $\beta \in [0, 1)$ let

$$(44) \quad 0 < \rho < \omega(\theta|\beta) - 1,$$

there are infinitely many positive integers n ,

$$(45) \quad n\theta = m_1 + \beta + b_n, \quad |b_n| \leq \frac{1}{n^\rho}$$

and some $\lambda \in \mathbb{S}(-\tau)$ with

$$(46) \quad -n\tau = m + \lambda, \quad m = \lfloor -n\tau \rfloor.$$

For each such λ, β, n, m , we have

$$(47) \quad L_n^{(\alpha)}(x_n(z, s); q) = \frac{(-zq^\alpha)^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}}{(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}} \times \left\{ \Theta \left(-zq^\alpha q^{X(m)+\lambda} e^{-2\beta\pi i} \mid q \right) + r_{ql}(n|5) \right\}$$

with

$$(48) \quad |r_{ql}(n|5)| \leq \frac{96(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q})}{(1-q)^4(q; q)_\infty} \times \left\{ (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} + \frac{\log^2 n}{n^\rho} \right\}$$

and

$$(49) \quad \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor$$

for n sufficiently large.

- (6) When $-2 < \tau < 0$, τ is irrational and θ is rational, Given a real number $\beta \in [0, 1)$ let

$$(50) \quad 0 < \rho < \omega(\theta|\beta) - 1,$$

there are infinitely many positive integers n ,

$$(51) \quad -n\tau = m + \beta + a_n, \quad |a_n| \leq \frac{1}{n^\rho}$$

and some $\lambda \in \mathbb{S}(\theta)$ with

$$(52) \quad n\theta = m_1 + \lambda, \quad \lambda = \{n\theta\}.$$

For each such λ, β, n, m , we have

$$(53) \quad \begin{aligned} L_n^{(\alpha)}(x_n(z, s); q) &= \frac{(-zq^\alpha)^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}}{(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}} \\ &\times \left\{ \Theta \left(-zq^\alpha q^{\chi(m) + \beta} e^{-2\lambda\pi i} \mid q \right) + r_{qt}(n|6) \right\} \end{aligned}$$

with

$$(54) \quad \begin{aligned} |r_{qt}(n|6)| &\leq \frac{96(-q^2; q)_\infty^3 \Theta(|z|q^\alpha \mid \sqrt{q})}{(1-q)^4 (q; q)_\infty} \\ &\times \left\{ (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} + \frac{\log^2 n}{n^\rho} \right\} \end{aligned}$$

and

$$(55) \quad \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor$$

for n sufficiently large.

- (7) When $-2 < \tau < 0$, both τ and θ are irrational. Assume that there exist two real numbers $\beta_1, \beta_2 \in [0, 1)$ with $\omega(-\tau, \theta | \beta_1, \beta_2) > 1$. Then for any

$$(56) \quad 0 < \rho < \omega(-\tau, \theta | \beta_1, \beta_2) - 1$$

there are infinitely many positive integers n such that

$$(57) \quad -\tau n = m + \beta_1 + a_n, \quad |a_n| < \frac{1}{n^\rho}$$

and

$$(58) \quad n\theta = m_1 + \beta_2 + b_n, \quad |b_n| < \frac{1}{n^\rho}.$$

For each such β_1, β_2, n, m , we have

$$(59) \quad \begin{aligned} L_n^{(\alpha)}(x_n(z, s); q) &= \frac{(-zq^\alpha)^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}}{(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}} \\ &\times \left\{ \Theta \left(-zq^\alpha q^{\chi(m) + \beta_1} e^{-2\beta_2\pi i} \mid q \right) + r_{qt}(n|7) \right\} \end{aligned}$$

with

$$(60) \quad \begin{aligned} |r_{qt}(n|7)| &\leq \frac{96(-q^2; q)_\infty^3 \Theta(|z|q^\alpha \mid \sqrt{q})}{(1-q)^4 (q; q)_\infty} \\ &\times \left\{ (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} + \frac{\log^2 n}{n^\rho} \right\} \end{aligned}$$

and

$$(61) \quad \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor$$

for n sufficiently large.

Remark 3.2. In the cases when only τ or θ is irrational, say τ , if it is an algebraic number of degree l , then we could take the corresponding $\beta = 0$, and the order of the error term is no better than $\mathcal{O}(n^{1-l})$, when this number is a Liouville number, however, the order of the error term is better than any $\mathcal{O}(n^{-k})$. It is clear that the scaling (12) with $\tau \leq -2$ won't give us anything interesting.

4. q -LAGUERRE POLYNOMIALS

In this section, we have used the following inequalities

$$(62) \quad 0 < (a; q)_n \leq 1, \quad (-b, q)_n \geq 1$$

for $0 \leq a < 1$, $b \geq 0$ and for $n \in \mathbb{N}$ or $n = \infty$. In the proofs 4.1–4.3 we will use the inequalities

$$(63) \quad 0 < \frac{(q; q)_\infty}{(q; q)_{n-k}} \frac{(q^{\alpha+1}; q)_n}{(q^{\alpha+1}; q)_{n-k}} \leq 1$$

for $0 \leq k \leq n$ and

$$(64) \quad \left| \frac{(q; q)_\infty}{(q; q)_{n-k}} \frac{(q^{\alpha+1}; q)_n}{(q^{\alpha+1}; q)_{n-k}} - 1 \right| \leq \frac{7(-q^2; q)_\infty^2 q^{n/2}}{(1-q)^3 (q; q)_\infty}$$

for $0 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$ and $\alpha > -1$. This can be seen from

$$(65) \quad \begin{aligned} \frac{(q; q)_\infty}{(q; q)_{n-k}} \frac{(q^{\alpha+1}; q)_n}{(q^{\alpha+1}; q)_{n-k}} &= \{R_1(q; n-k) + 1\} \{R_2(q^{\alpha+1}; n) + 1\} \\ &\times \{R_1(q^{\alpha+1}; n-k) + 1\} \end{aligned}$$

and Lemma 2.1.

In the proofs 4.4–4.7, we have

$$(66) \quad -2 < \tau < 0.$$

and

$$(67) \quad 2 \ll \nu_n \leq n \min \left\{ \frac{2+\tau}{8}, \frac{-\tau}{8} \right\} < \frac{n}{4}$$

for n sufficiently large. Assume that

$$(68) \quad -\tau n = m + c_n, \quad 0 \leq c_n < 1, \quad 0 < m = \lfloor -\tau n \rfloor,$$

$$(69) \quad n\theta = m_1 + d_n, \quad 0 \leq d_n < 1, \quad m_1 = \lfloor n\theta \rfloor.$$

Then

$$(70) \quad \begin{aligned} \frac{L_n^{(\alpha)}(x_n(z, s); q)}{(-zq^\alpha)^n q^{n^2(1-s)}} &= \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{q^{k^2} e^{2nk\theta\pi i}}{(q^{\alpha+1}; q)_{n-k}} \left(-\frac{q^{\tau n}}{zq^\alpha} \right)^k \\ &= \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=0}^{\lfloor m/2 \rfloor} \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{q^{k^2} e^{2nk\theta\pi i}}{(q^{\alpha+1}; q)_{n-k}} \left(-\frac{q^{\tau n}}{zq^\alpha} \right)^k \\ &\quad + \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=\lfloor m/2 \rfloor + 1}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{q^{k^2} e^{2nk\theta\pi i}}{(q^{\alpha+1}; q)_{n-k}} \left(-\frac{q^{\tau n}}{zq^\alpha} \right)^k \\ &= s_1 + s_2. \end{aligned}$$

We reverse the summation order in s_1 to obtain

$$(71) \quad \frac{s_1(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \sum_{k=0}^{\lfloor m/2 \rfloor} q^{k^2} \left(-zq^\alpha q^{\chi(m) + c_n} e^{-2\pi i d_n} \right)^k e(k, n)$$

with

$$(72) \quad e(k, n) = \frac{(q; q)_\infty^2 (q^{\alpha+1}; q)_n}{(q; q)_{\lfloor m/2 \rfloor - k} (q; q)_{n - \lfloor m/2 \rfloor + k} (q^{\alpha+1}; q)_{n - \lfloor m/2 \rfloor + k}}.$$

Since $0 < q < 1$, $\alpha > -1$, it is clear that

$$(73) \quad |e(k, n)| \leq 1$$

for $0 \leq k \leq \lfloor \frac{m}{2} \rfloor$. Observe that

$$(74) \quad \begin{aligned} e(k, n) &= \{R_2(q^{\alpha+1}; n) + 1\} \left\{R_1(q; \lfloor \frac{m}{2} \rfloor - k) + 1\right\} \\ &\times \left\{R_1(q; n - \lfloor \frac{m}{2} \rfloor + k) + 1\right\} \left\{R_1(q^{\alpha+1}; n - \lfloor \frac{m}{2} \rfloor + k) + 1\right\}. \end{aligned}$$

By Lemma 2.1, we get

$$(75) \quad |e(k, n) - 1| \leq \frac{15(-q^2; q)_\infty^3 q^{\nu_n+1}}{(1-q)^4 (q; q)_\infty}$$

for $0 \leq k \leq \nu_n - 1$.

In sum s_2 we shift summation from k to $k + \lfloor \frac{m}{2} \rfloor$ to get

$$(76) \quad \frac{s_2(q; q)_\infty^2 (-z q^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \sum_{k=1}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-c_n} e^{2\pi i d_n} \right)^k f(k, n)$$

with

$$(77) \quad f(k, n) = \frac{(q; q)_\infty^2 (q^{\alpha+1}; q)_n}{(q; q)_{\lfloor m/2 \rfloor + k} (q; q)_{n-\lfloor m/2 \rfloor - k} (q^{\alpha+1}; q)_{n-\lfloor m/2 \rfloor - k}}$$

for $1 \leq k \leq n - \lfloor \frac{m}{2} \rfloor$. Hence

$$(78) \quad |f(k, n)| \leq 1$$

for $1 \leq k \leq n - \lfloor \frac{m}{2} \rfloor$. Apply Lemma 2.1 to

$$(79) \quad \begin{aligned} f(k, n) &= \{R_2(q^{\alpha+1}; n) + 1\} \times \left\{R_1(q; \lfloor \frac{m}{2} \rfloor + k) + 1\right\} \\ &\times \left\{R_1(q; n - \lfloor \frac{m}{2} \rfloor - k) + 1\right\} \left\{R_1(q^{\alpha+1}; n - \lfloor \frac{m}{2} \rfloor - k) + 1\right\}, \end{aligned}$$

we obtain

$$(80) \quad |f(k, n) - 1| \leq \frac{15(-q^2; q)_\infty^3 q^{\nu_n+1}}{(1-q)^4 (q; q)_\infty}$$

for $1 \leq k \leq \nu_n - 1$.

4.1. Proof for the case $\tau > 0$.

Proof. From (13) one has

$$(81) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-z q^\alpha)^n q^{n^2(1-s)}} = 1 + r_{ql}(n|1),$$

and

$$(82) \quad r_{ql}(n|1) = \sum_{k=1}^n \frac{q^{k^2} e^{2nk\theta\pi i}}{(q; q)_k} \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \left(-\frac{q^{\tau n}}{z q^\alpha} \right)^k.$$

Thus

$$\begin{aligned}
 |r_{ql}(n|1)| &\leq \sum_{k=1}^n \frac{q^{k^2}}{(q; q)_k} \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \left(\frac{q^{\tau n}}{|z| q^\alpha} \right)^k \\
 &\leq \sum_{k=1}^n \frac{q^{k^2}}{(q; q)_k} \left(\frac{q^{\tau n}}{|z| q^\alpha} \right)^k \\
 &\leq \sum_{k=1}^\infty \frac{q^{k^2}}{(q; q)_k} \left(\frac{q^{\tau n}}{|z| q^\alpha} \right)^k \\
 (83) \quad &\leq \frac{q^{1-\alpha} B_q(q^{2-\alpha} |z|^{-1})}{(1-q)|z|} q^{\tau n}.
 \end{aligned}$$

□

4.2. Proof for the case $\tau = 0$ and θ rational.

Proof. We have

$$\begin{aligned}
 \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} &= \sum_{k=0}^n \frac{q^{k^2}}{(q; q)_k} \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \left(-\frac{e^{2\lambda\pi i}}{zq^\alpha} \right)^k \\
 &= \sum_{k=0}^\infty \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\lambda\pi i}}{zq^\alpha} \right)^k \\
 &\quad - \sum_{k=\lfloor n/2 \rfloor}^\infty \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\lambda\pi i}}{zq^\alpha} \right)^k \\
 &\quad + \sum_{k=0}^{\lfloor n/2 \rfloor - 1} \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\lambda\pi i}}{zq^\alpha} \right)^k \left\{ \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} - 1 \right\} \\
 &\quad + \sum_{k=\lfloor n/2 \rfloor}^n \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\lambda\pi i}}{zq^\alpha} \right)^k \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \\
 (84) \quad &= A_q \left(\frac{e^{2\lambda\pi i}}{zq^\alpha} \right) + s_1 + s_2 + s_3.
 \end{aligned}$$

Then

$$\begin{aligned}
 |s_1 + s_3| &\leq 2 \sum_{k=\lfloor n/2 \rfloor}^\infty \frac{q^{k^2}}{(q; q)_k} \left(\frac{1}{|z| q^\alpha} \right)^k \\
 (85) \quad &= 2 \frac{q^{n^2/4} B_q(|z|^{-1} q^{-\alpha})}{(q; q)_\infty (|z| q^\alpha)^{\lfloor n/2 \rfloor}},
 \end{aligned}$$

and

$$(86) \quad |s_2| \leq \frac{7(-q^2; q)_\infty^2 B_q(|z|^{-1} q^{-\alpha}) q^{n/2}}{(1-q)^3 (q; q)_\infty}.$$

Therefore

$$(87) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} = A_q \left(\frac{e^{2\lambda\pi i}}{zq^\alpha} \right) + r_{ql}(n|2)$$

with

$$(88) \quad r_{ql}(n|2) = s_1 + s_2 + s_3$$

and

$$(89) \quad |r_{ql}(n|2)| \leq \frac{7(-q^2; q)_\infty^2 B_q(|z|^{-1} q^{-\alpha})}{(1-q)^3 (q; q)_\infty} \left\{ q^{n/2} + \frac{q^{n^2/4}}{(|z|q^\alpha)^{\lfloor n/2 \rfloor}} \right\}.$$

□

4.3. Proof for the case $\tau = 0$ and θ irrational.

Proof. Assume that

$$(90) \quad \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor,$$

we have

$$\begin{aligned} \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-zq^\alpha)^n q^{n^2(1-s)}} &= \sum_{k=0}^n \frac{q^{k^2} e^{2\pi i k \gamma_n}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \\ &= \sum_{k=0}^\infty \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \\ &\quad - \sum_{k=\nu_n}^\infty \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \\ &\quad + \sum_{k=0}^{\nu_n-1} \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \left\{ \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} - 1 \right\} \\ &\quad + \sum_{k=0}^{\nu_n-1} \frac{q^{k^2}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \{e^{2\pi i k \gamma_n} - 1\} \\ &\quad + \sum_{k=\nu_n}^n \frac{q^{k^2} e^{2\pi i k \gamma_n}}{(q; q)_k} \left(-\frac{e^{2\pi i \beta}}{zq^\alpha} \right)^k \frac{(q^{\alpha+1}; q)_n (q; q)_\infty}{(q^{\alpha+1}; q)_{n-k} (q; q)_{n-k}} \\ (91) \quad &= A_q \left(\frac{e^{2\pi i \beta}}{zq^\alpha} \right) + s_1 + s_2 + s_3 + s_4. \end{aligned}$$

Then,

$$\begin{aligned} |s_1 + s_4| &\leq 2 \sum_{k=\nu_n}^\infty \frac{q^{k^2} |z|^{-k}}{(q; q)_k q^{k\alpha}} \\ (92) \quad &\leq \frac{2B_q(|z|^{-1} q^{-\alpha}) q^{\nu_n^2}}{(q; q)_\infty (|z|q^\alpha)^{\nu_n}}. \end{aligned}$$

Clearly, we have

$$(93) \quad \nu_n \ll n^{\min(1, \rho)} / 8$$

and

$$(94) \quad q^{n/2} \ll \frac{\nu_n}{n^\rho}$$

for sufficiently large n . Hence

$$(95) \quad |s_2| \leq \frac{7(-q^2; q)_\infty^2 B_q(|z|^{-1} q^{-\alpha}) \log^2 n}{(1-q)^3 (q; q)_\infty n^\rho}.$$

It is clear that for large n ,

$$(96) \quad |e^{2\pi i k \gamma_n} - 1| \leq \frac{2\pi \nu_n}{n^\rho} e^{2\pi \nu_n / n^\rho} \leq \frac{24 \log^2 n}{n^\rho},$$

and

$$(97) \quad |s_3| \leq \frac{24 B_q(|z|^{-1} q^{-\alpha}) \log^2 n}{n^\rho}.$$

Let

$$(98) \quad e_{ql}(n|3) = s_1 + s_2 + s_3 + s_4,$$

then

$$(99) \quad \frac{L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty}{(-z q^\alpha)^n q^{n^2(1-s)}} = A_q \left(\frac{e^{2\pi i \beta}}{z q^\alpha} \right) + e_{ql}(n|3)$$

with

$$(100) \quad |e_{ql}(n|3)| \leq \frac{48(-q^2; q)_\infty^2 B_q(|z|^{-1} q^{-\alpha})}{(1-q)^3 (q; q)_\infty} \left\{ \frac{\log^2 n}{n^\rho} + \frac{q^{\nu_n^2}}{(|z| q^\alpha)^{\nu_n}} \right\}$$

for n sufficiently large. □

4.4. Proof for the case $-2 < \tau < 0$ rational and θ rational.

Proof. Assume that

$$(101) \quad \nu_n = \min \left\{ \left\lfloor \frac{(2+\tau)n}{8} \right\rfloor, \left\lfloor \frac{-\tau n}{8} \right\rfloor \right\}.$$

From (71) we get

$$(102) \quad \begin{aligned} \frac{s_1(-z q^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor} (q; q)_\infty^2}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=0}^{\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k \\ &- \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k \\ &+ \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k (e(k, n) - 1) \\ &+ \sum_{k=\nu_n}^{\lfloor m/2 \rfloor} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k e(k, n) \\ &= \sum_{k=0}^{\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k + s_{11} + s_{12} + s_{13}. \end{aligned}$$

Then,

$$\begin{aligned}
 |s_{11} + s_{12}| &\leq 2 \sum_{k=\nu_n}^{\infty} q^{k^2} \left(|z| q^{\alpha} q^{\chi(m)+\lambda} \right)^k \\
 &\leq 2 \left(|z| q^{\alpha} \right)^{\nu_n} q^{\nu_n^2} \sum_{k=0}^{\infty} q^{k^2} \left(|z| q^{\alpha} q^{\chi(m)+\lambda} \right)^k \\
 (103) \quad &\leq 2\Theta \left(|z| q^{\alpha} \mid \sqrt{q} \right) \left(|z| q^{\alpha} \right)^{\nu_n} q^{\nu_n^2}
 \end{aligned}$$

and

$$\begin{aligned}
 |s_{12}| &\leq \frac{15(-q^2; q)_{\infty}^3 q^{\nu_n}}{(1-q)^4 (q; q)_{\infty}} \sum_{k=0}^{\infty} q^{k^2} \left(|z| q^{\alpha} q^{\chi(m)+\lambda} \right)^k \\
 (104) \quad &\leq \frac{15(-q^2; q)_{\infty}^3 \Theta \left(|z| q^{\alpha} \mid \sqrt{q} \right) q^{\nu_n}}{(1-q)^4 (q; q)_{\infty}}.
 \end{aligned}$$

Let

$$(105) \quad r_1(n) = s_{11} + s_{12} + s_{13},$$

then,

$$(106) \quad \frac{s_1(q; q)_{\infty}^2 (-z q^{\alpha} e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \sum_{k=0}^{\infty} q^{k^2} \left(-z q^{\alpha} q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k + r_1(n)$$

$$(107) \quad |r_1(n)| \leq \frac{15(-q^2; q)_{\infty}^3 \Theta \left(|z| q^{\alpha} \mid \sqrt{q} \right)}{(1-q)^4 (q; q)_{\infty}} \left\{ \left(|z| q^{\alpha} \right)^{\nu_n} q^{\nu_n^2} + q^{\nu_n} \right\}.$$

From (76) we obtain,

$$\begin{aligned}
 \frac{s_2(-z q^{\alpha} e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor} (q; q)_{\infty}^2}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=1}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\pi i \lambda_1} \right)^k \\
 &- \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\pi i \lambda_1} \right)^k \\
 &+ \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\pi i \lambda_1} \right)^k (f(k, n) - 1) \\
 &+ \sum_{k=\nu_n}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\pi i \lambda_1} \right)^k f(k, n) \\
 (108) \quad &= \sum_{k=-1}^{-\infty} q^{k^2} \left(-z q^{\alpha} q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k + s_{21} + s_{22} + s_{23},
 \end{aligned}$$

then,

$$\begin{aligned}
 |s_{21} + s_{23}| &\leq 2 \sum_{k=\nu_n}^{\infty} q^{k^2-2k} \left(|z|^{-1} q^{-\alpha} \right)^k \\
 &\leq 2 \frac{q^{\nu_n^2-2\nu_n-2} \sum_{k=0}^{\infty} q^{k^2/2} |z|^{-k} q^{-\alpha k}}{|z|^{\nu_n} q^{\alpha \nu_n}} \\
 &\leq 2 \frac{q^{\nu_n^2/2} \sum_{k=0}^{\infty} q^{k^2/2} |z|^{-k} q^{-\alpha k}}{|z|^{\nu_n} q^{\alpha \nu_n}} \\
 (109) \quad &\leq 2 \frac{q^{\nu_n^2/2} \Theta(|z| q^{\alpha} | \sqrt{q})}{(|z| q^{\alpha})^{\nu_n}}
 \end{aligned}$$

for n sufficiently large. For the sum s_{22} , we have

$$\begin{aligned}
 |s_{22}| &\leq \frac{15(-q^2; q)_{\infty}^3 q^{\nu_n+1}}{(1-q)^4 (q; q)_{\infty}} \sum_{k=1}^{\infty} q^{k^2/2+k^2/2-2k} |z|^{-k} q^{-\alpha k} \\
 &\leq \frac{15(-q^2; q)_{\infty}^3 q^{\nu_n-1}}{(1-q)^4 (q; q)_{\infty}} \sum_{k=1}^{\infty} q^{k^2/2} |z|^{-k} q^{-\alpha k} \\
 (110) \quad &\leq \frac{15(-q^2; q)_{\infty}^3 \Theta(|z| q^{\alpha} | \sqrt{q}) q^{\nu_n/2}}{(1-q)^4 (q; q)_{\infty}}.
 \end{aligned}$$

Hence

$$(111) \quad \frac{s_2(-z q^{\alpha} e^{-2n\theta\pi i})_{\lfloor m/2 \rfloor} (q; q)_{\infty}^2}{q^{\lfloor m/2 \rfloor} (\tau n + \lfloor m/2 \rfloor)} = \sum_{k=-1}^{\infty} q^{k^2} \left(-z q^{\alpha} q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \right)^k + r_2(n)$$

with

$$(112) \quad r_2(n) = s_{21} + s_{22} + s_{23}$$

and

$$(113) \quad |r_2(n)| \leq \frac{15(-q^2; q)_{\infty}^3 \Theta(|z| q^{\alpha} | \sqrt{q})}{(1-q)^4 (q; q)_{\infty}} \left\{ q^{\nu_n/2} + \frac{q^{\nu_n^2/2}}{(|z| q^{\alpha})^{\nu_n}} \right\}.$$

Thus

$$(114) \quad \frac{(-z q^{\alpha} e^{-2n\theta\pi i})_{\lfloor m/2 \rfloor} L_n^{(\alpha)}(x_n(z, s); q) (q; q)_{\infty}^2}{(-z q^{\alpha})^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \Theta \left(-z q^{\alpha} q^{\chi(m)+\lambda} e^{-2\pi i \lambda_1} \mid q \right) + r_{ql}(n|4)$$

with

$$(115) \quad |r_{ql}(n|4)| \leq \frac{15(-q^2; q)_{\infty}^3 \Theta(|z| q^{\alpha} | \sqrt{q})}{(1-q)^4 (q; q)_{\infty}} \left\{ q^{\nu_n/2} + (|z| q^{\alpha})^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z| q^{\alpha})^{\nu_n}} \right\}$$

for n sufficiently large.

□

4.5. Proof for the case $-2 < \tau < 0$ rational and θ irrational.

Proof. The existence of λ with infinitely many positive integers n satisfying (45) and (46) is guaranteed by the fact that $\mathbb{S}(-\tau)$ is a finite set.

It is clear that for sufficiently large n , we have

$$(116) \quad 2 \ll \nu_n = \left\lfloor \frac{q^4 \log^2 n}{1 + \log q^{-1}} \right\rfloor \ll \frac{n^\rho}{8}, \quad q^{\nu_n} \ll \frac{\nu_n}{n^\rho}.$$

From (71) to get

$$\begin{aligned} \frac{s_1(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=0}^{\lfloor m/2 \rfloor} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k e^{-2k\pi i b_n} e(k, n) \\ &= \sum_{k=0}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k \\ &\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k \\ &\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k \{e^{-2k\pi i b_n} - 1\} \\ &\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k e^{-2k\pi i b_n} \{e(k, n) - 1\} \\ &\quad + \sum_{k=\nu_n}^{\lfloor m/2 \rfloor} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k e^{-2k\pi i b_n} e(k, n) \\ (117) \quad &= \sum_{k=0}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i\beta} \right)^k + s_{11} + s_{12} + s_{13} + s_{14}, \end{aligned}$$

then

$$\begin{aligned} |s_{11} + s_{14}| &\leq 2 \sum_{k=\nu_n}^{\infty} q^{k^2} |z|^k q^{\alpha k} \\ &\leq 2 (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} \sum_{k=0}^{\infty} q^{k^2} |z|^k q^{\alpha k} \\ (118) \quad &\leq 2\Theta(|z|q^\alpha | \sqrt{q}) (|z|q^\alpha)^{\nu_n} q^{\nu_n^2}. \end{aligned}$$

For sufficiently large n ,

$$\begin{aligned} |s_{12}| &\leq \frac{24\nu_n}{n^\rho} \sum_{k=0}^{\infty} q^{k^2} |z|^k q^{\alpha k} \\ (119) \quad &\leq \frac{24\Theta(|z|q^\alpha | \sqrt{q}) \log^2 n}{n^\rho}, \end{aligned}$$

and

$$\begin{aligned}
 |s_{13}| &\leq \frac{15(-q^2; q)_\infty^3 q^{\nu_n+1}}{(1-q)^4 (q; q)_\infty} \sum_{k=0}^{\infty} q^{k^2} |z|^k q^{\alpha k} \\
 (120) \quad &\leq \frac{15(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q}) \log^2 n}{(1-q)^4 (q; q)_\infty n^\rho}.
 \end{aligned}$$

Hence,

$$(121) \quad \frac{s_1(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \sum_{k=0}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\pi i \beta} \right)^k + r_1(n)$$

with

$$(122) \quad r_1(n) = s_{11} + s_{12} + s_{13} + s_{14}$$

and

$$(123) \quad |r_1(n)| \leq \frac{48(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q})}{(1-q)^4 (q; q)_\infty} \left\{ (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{\log^2 n}{n^\rho} \right\}$$

for n sufficiently large.

From (76) we obtain

$$\begin{aligned}
 \frac{s_2(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=1}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k e^{2k\pi i b_n} f(k, n) \\
 &= \sum_{k=1}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k \\
 &\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k \\
 &\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k \{e^{2k\pi i b_n} - 1\} \\
 &\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k e^{2k\pi i b_n} \{f(k, n) - 1\} \\
 &\quad + \sum_{k=\nu_n}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\lambda} e^{2\beta\pi i} \right)^k e^{2k\pi i b_n} f(k, n) \\
 (124) \quad &= \sum_{k=-1}^{-\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\beta\pi i} \right)^k + s_{21} + s_{22} + s_{23} + s_{24},
 \end{aligned}$$

then

$$\begin{aligned}
 |s_{21} + s_{24}| &\leq 2 \sum_{k=\nu_n}^{\infty} q^{k^2-2k} |z|^{-k} q^{-\alpha k} \\
 &\leq 2 (|z|q^\alpha)^{-\nu_n} q^{\nu_n^2-2\nu_n} \sum_{k=0}^{\infty} q^{k^2/2} |z|^{-k} q^{-\alpha k} \\
 (125) \quad &\leq \frac{2\Theta(|z|q^\alpha | \sqrt{q}) q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}}
 \end{aligned}$$

and

$$\begin{aligned}
 |s_{22}| &\leq \frac{24\nu_n}{n^\rho} \sum_{k=0}^{\infty} q^{k^2/2+k^2/2-2k} |z|^{-k} q^{-\alpha k} \\
 (126) \qquad &\leq \frac{24\Theta(|z|q^\alpha | \sqrt{q}) \log^2 n}{n^\rho}
 \end{aligned}$$

and

$$\begin{aligned}
 |s_{23}| &\leq \frac{15(-q^2; q)_\infty^3 q^{\nu_n+1}}{(1-q)^4(q; q)_\infty} \sum_{k=0}^{\infty} q^{k^2/2+k^2/2-2k} |z|^{-k} q^{-\alpha k} \\
 &\leq \frac{15(-q^2; q)_\infty^3 q^{\nu_n}}{(1-q)^4(q; q)_\infty} \sum_{k=0}^{\infty} q^{k^2/2} |z|^{-k} q^{-\alpha k} \\
 (127) \qquad &\leq \frac{15(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q}) \log^2 n}{(1-q)^4(q; q)_\infty} \frac{1}{n^\rho}.
 \end{aligned}$$

for n sufficiently large. Therefore

$$(128) \quad \frac{s_2(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \sum_{k=-1}^{-\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\beta\pi i} \right)^k + r_2(n)$$

with

$$(129) \qquad r_2(n) = s_{21} + s_{22} + s_{23} + s_{24}$$

and

$$(130) \qquad |r_2(n)| \leq \frac{48(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q})}{(1-q)^4(q; q)_\infty} \left\{ \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} + \frac{\log^2 n}{n^\rho} \right\}$$

for n sufficiently large. Combine (121) and (128) we get

$$(131) \quad \frac{(-zq^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor} L_n^{(\alpha)}(x_n(z, s); q)(q; q)_\infty^2}{(-zq^\alpha)^n q^{n^2(1-s) + \lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} = \Theta \left(-zq^\alpha q^{\chi(m)+\lambda} e^{-2\beta\pi i} \mid q \right) + r_{ql}(n|5)$$

with

$$(132) \qquad r_{ql}(n|5) = r_1(n) + r_2(n)$$

and

$$(133) \qquad |r_{ql}(n|5)| \leq \frac{96(-q^2; q)_\infty^3 \Theta(|z|q^\alpha | \sqrt{q})}{(1-q)^4(q; q)_\infty} \left\{ (|z|q^\alpha)^{\nu_n} q^{\nu_n^2} + \frac{q^{\nu_n^2/2}}{(|z|q^\alpha)^{\nu_n}} + \frac{\log^2 n}{n^\rho} \right\}$$

for n sufficiently large. □

4.6. Proof for the case $-2 < \tau < 0$ irrational and θ rational.

Proof. This proof is very similar to the proof for τ rational and θ irrational . We just observe that

$$\begin{aligned}
\frac{s_1(q; q)_\infty^2 (-z q^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=0}^{\lfloor m/2 \rfloor} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k q^{ka_n} e(k, n) \\
&= \sum_{k=0}^{\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k \\
&\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k \\
&\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k \{q^{ka_n} - 1\} \\
&\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k q^{ka_n} \{e(k, n) - 1\} \\
(134) \quad &+ \sum_{k=\nu_n}^{\lfloor m/2 \rfloor} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k q^{ka_n} e(k, n),
\end{aligned}$$

and

$$\begin{aligned}
\frac{s_2(q; q)_\infty^2 (-z q^\alpha e^{-2n\theta\pi i})^{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=1}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta} e^{2\lambda\pi i} \right)^k q^{-ka_n} f(k, n) \\
&= \sum_{k=-1}^{-\infty} q^{k^2} \left(-z q^\alpha q^{\chi(m)+\beta} e^{-2\pi i \lambda} \right)^k \\
&\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta} e^{2\lambda\pi i} \right)^k \\
&\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta} e^{2\lambda\pi i} \right)^k \{q^{-ka_n} - 1\} \\
&\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta} e^{2\lambda\pi i} \right)^k q^{-ka_n} \{f(k, n) - 1\} \\
(135) \quad &+ \sum_{k=\nu_n}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta} e^{2\lambda\pi i} \right)^k q^{-ka_n} f(k, n).
\end{aligned}$$

□

4.7. Proof for the case $-2 < \tau < 0$ irrational and θ irrational.

Proof. In this case we have

$$\begin{aligned}
\frac{s_1(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})_{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=0}^{\lfloor m/2 \rfloor} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k e^{-2k\pi i b_n} q^{ka_n} e(k, n) \\
&= \sum_{k=0}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k \\
&\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k \\
&\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k \{q^{ka_n} - 1\} \\
&\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k \{e^{-2k\pi i b_n} - 1\} q^{ka_n} \\
&\quad + \sum_{k=0}^{\nu_n-1} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k e^{-2k\pi i b_n} q^{ka_n} \{e(k, n) - 1\} \\
&\quad + \sum_{k=\nu_n}^{\lfloor m/2 \rfloor} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\pi i\beta_2} \right)^k e^{-2k\pi i b_n} q^{ka_n} e(k, n),
\end{aligned}$$

and

$$\begin{aligned}
\frac{s_2(q; q)_\infty^2 (-zq^\alpha e^{-2n\theta\pi i})_{\lfloor m/2 \rfloor}}{q^{\lfloor m/2 \rfloor (\tau n + \lfloor m/2 \rfloor)}} &= \sum_{k=1}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k e^{2k\pi i b_n} q^{-ka_n} f(k, n) \\
&= \sum_{k=-1}^{-\infty} q^{k^2} \left(-zq^\alpha q^{\chi(m)+\beta_1} e^{-2\beta_2\pi i} \right)^k \\
&\quad - \sum_{k=\nu_n}^{\infty} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k \\
&\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k \{e^{2k\pi i b_n} - 1\} \\
&\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k e^{2k\pi i b_n} \{q^{-ka_n} - 1\} \\
&\quad + \sum_{k=1}^{\nu_n-1} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k e^{2k\pi i b_n} q^{-ka_n} \{f(k, n) - 1\} \\
&\quad + \sum_{k=\nu_n}^{n-\lfloor m/2 \rfloor} q^{k^2} \left(-z^{-1} q^{-\alpha} q^{-\chi(m)-\beta_1} e^{2\beta_2\pi i} \right)^k e^{2k\pi i b_n} q^{-ka_n} f(k, n).
\end{aligned}$$

To finish the proof, we have to estimate the sums above, but they are very similar to what we have done in the other cases, we won't repeat these details here. $\square$

REFERENCES

- [1] N. I. Akhiezer, The Classical Moment Problem and Some Related Questions in Analysis, English translation, Oliver and Boyd, Edinburgh, 1965.
- [2] Tom Apostol, Modular Functions and Dirichlet Series in Number Theory, Second Edition, Springer-Verlag, New York, 1990.
- [3] G. E. Andrews, q -series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra, CBMS Regional Conference Series, number 66, American Mathematical Society, Providence, R.I. 1986.
- [4] G. E. Andrews, Ramanujan's "Lost" Note book VIII: The entire Rogers-Ramanujan function, *Advances in Math.* 191 (2005), 393–407.
- [5] G. E. Andrews, Ramanujan's "Lost" Note book IX: The entire Rogers-Ramanujan function, *Advances in Math.* 191 (2005), 408–422.
- [6] G. E. Andrews, R. A. Askey, and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.
- [7] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach*, American Mathematical Society, Providence, 2000.
- [8] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.* 52 (1999), 1491–1552.
- [9] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition Cambridge University Press, Cambridge, 2004.
- [10] W. K. Hayman, On the zeros of a q -Bessel function, *Contemporary Mathematics*, volume 382, American Mathematical Society, Providence, 2005, 205–216.
- [11] Hua Loo Keng, *Introduction to Number Theory*, Springer-Verlag, Berlin Heidelberg New York, 1982.
- [12] M. E. H. Ismail, Asymptotics of q -orthogonal polynomials and a q -Airy function, *Internat. Math. Res. Notices* 2005 No 18 (2005), 1063–1088.
- [13] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one Variable*, Cambridge University Press, Cambridge, 2005.
- [14] M. E. H. Ismail and X. Li, Bounds for extreme zeros of orthogonal polynomials, *Proc. Amer. Math. Soc.* 115 (1992), 131–140.
- [15] M. E. H. Ismail and D. R. Masson, q -Hermite polynomials, biorthogonal rational functions, *Trans. Amer. Math. Soc.* 346 (1994), 63–116.
- [16] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, *Advances in Math.*, (2007), to appear.
- [17] M. E. H. Ismail and R. Zhang, Scaled asymptotics for q -polynomials, *Comptes Rendus*, submitted.
- [18] M. E. H. Ismail and R. Zhang, Chaotic and Periodic Asymptotics for q -Orthogonal Polynomials, joint with Mourad E.H. Ismail, *International Mathematics Research Notices*, accepted.
- [19] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, Y. Yamada, Hypergeometric solutions to the q -Painlevé equations, *Internat. Math. Res. Notices* 47 (2004), 2497–2521.
- [20] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogues, *Reports of the Faculty of Technical Mathematics and Informatics* no. 98-17, Delft University of Technology, Delft, 1998.
- [21] M. L. Mehta, *Random Matrices*, third edition, Elsevier, Amsterdam, 2004.
- [22] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers* (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.
- [23] E. B. Saff and V. Totik, *Logarithmic Potentials With External Fields*, Springer-Verlag, New York, 1997.
- [24] G. Szegő, *Orthogonal Polynomials*, Fourth Edition, Amer. Math. Soc., Providence, 1975.
- [25] R. Wong, *Asymptotic Approximations of Integrals*, Academic Press, Boston, 1989.
- [26] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, fourth edition, Cambridge University Press, Cambridge, 1927.
- [27] Wikipedia.org, http://en.wikipedia.org/wiki/Liouville_number.
- [28] Ruiming Zhang, Planchère-Rotach Asymptotics for Stieltjes-Wigert Orthogonal Polynomials with Complex Scaling.

- [29] Ruiming Zhang, Planchere-Rotach Asymptotics for Ismail-Masson Orthogonal Polynomials with Complex Scaling.

Current address: School of Mathematics, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.

E-mail address: ruimingzhang@yahoo.com